



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

No. 6.

Telegram of the sanitary inspector of Djiddah.

DJIDDAH, April 25, 1900.

Pilgrim arrivals ended; 34,100. Health perfect.

DR. POMPOURAS.

No. 7.

Telegram of sanitary inspector of Camaran.

CAMARAN, March 30, 1900.

Sailing vessel *Hassim*, English flag, having stopped twenty-five days at Aden, arrived March 13 with 33 passengers from Yemen. One sailor presented fever, bubo, and prostration. Patient rigorously isolated; bed clothing and wearing apparel purchased at Aden burned. Proceeding with unloading cargo taken at Aden. Quarantine to last fifteen days.

XANTHROPAULIDES.

No. 8.

Circular telegram to health officers on the Turco-Persian frontier, dated April 21, 1900.

By decision of the superior council of health access to the Ottoman frontier comprised between Revendanz and Haneguine is prohibited, except by the three points of Revendanz, Pendjovine, and Haneguine, where travelers shall be subject to rigorous medical visit. Those travelers who present symptoms suspect disease shall not be received in Ottoman territory.

DR. COZZONIS,
Inspector-General.

FOREIGN STATISTICAL REPORTS.

AUSTRALIA—*New South Wales—Sydney*.—Month of February, 1900. Estimated population, 438,300. Total number of deaths, 456, including diphtheria, 2; enteric fever, 17; scarlet fever, 1; phthisis pulmonalis, 37, and 1 from bubonic plague.

FRANCE—*Bordeaux*.—Month of March, 1900. Estimated population, 286,000. Total number of deaths, 569, including diphtheria, 12; enteric fever, 4; measles, 7, and 3 from whooping cough.

Nice.—Month of March, 1900. Estimated population, 108,227. Total number of deaths, 259, including diphtheria, 2; enteric fever, 4; measles, 4; whooping cough, 3, and 32 from tuberculosis.

Rouen.—Month of March, 1900. Estimated population, 112,657. Total number of deaths, 329, including enteric fever, 5; measles, 1; whooping cough, 2; smallpox, 1, and 56 from tuberculosis.

St. Etienne.—Two weeks ended March 15, 1900. Estimated population, 135,784. Total number of deaths, 168, including diphtheria, 1; measles, 1, and 29 from phthisis pulmonalis.

GREAT BRITAIN—*England and Wales*.—The deaths registered in 33 great towns in England and Wales during the week ended April 7 1900, correspond to an annual rate of 23.0 a thousand of the aggregate population, which is estimated at 11,610,296. The highest rate was recorded in Wolverhampton, viz, 33.2, and the lowest in Derby, viz, 15.0.

London.—One thousand nine hundred and thirty-two deaths were registered during the week, including measles, 62; scarlet fever, 7; diphtheria, 32; whooping cough, 50; enteric fever, 20; and diarrhea and

dysentery, 6. The deaths from all causes correspond to an annual rate of 22.0 a thousand. In Greater London 2,536 deaths were registered, corresponding to an annual rate of 19.9 a thousand of the population. In the "outer ring" the deaths included 12 from diphtheria, 5 from measles, 1 from scarlet fever, and 18 from whooping cough.

Ireland.—The average annual death rate represented by the deaths registered during the week ended April 7, 1900, in the 22 principal town districts of Ireland was 26.4 a thousand of the population, which is estimated at 1,062,188. The lowest rate was recorded in Londonderry, viz, 6.3, and the highest in Armagh, viz, 64.1 a thousand. In Dublin and suburbs 194 deaths were registered, including enteric fever, 3; measles, 3, and 3 from whooping cough.

Scotland.—The deaths registered in 8 principal towns during the week ended April 7, 1900, correspond to an annual rate of 21.7 a thousand of the population, which is estimated at 1,606,935. The lowest mortality was recorded in Dundee, viz, 16.4, and the highest in Greenock, viz, 25.8 a thousand. The aggregate number of deaths registered from all causes was 672, including diphtheria, 6; measles, 14; scarlet fever, 4, and 15 from whooping cough.

NICARAGUA—*San Juan del Norte.*—Month of March, 1900. Estimated population, 1,156. Total number of deaths, 3. No contagious diseases.

NORFOLK ISLAND.—Month of January, 1900. Estimated population, 846. One death. No contagious diseases.

Month of February, 1900. No deaths. No contagious diseases.